

Product datasheet for **AP21233SU-N**

Ceruloplasmin (CP) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ID, IP
Recommended Dilution:	In precipitating techniques as Immuno-electrophoresis and single or Double Radial Immunodiffusion to identify the presence of alpha-2 Ceruloplasmin in Human plasma or other body fluids or to determine its concentration. The presence of non precipitating antibodies has not been assayed. This does not exclude the use of the antiserum in more sensitive non-precipitating antibody binding techniques if proper controls are included. <i>Recommended Dilutions:</i> Immuno-electrophoresis: Use 2 µl or equivalent against 120 µl antiserum. Double Radial Immunodiffusion (Ouchterlony): Use a rosette arrangement with 10 µl in a 3 mm diameter center well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Ceruloplasmin is isolated from pooled Human serum by sequential precipitation and purified by Ion Exchange Chromatography and Affinity Chromatography. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	The reactivity of the antiserum is restricted to alpha-2 Ceruloplasmin. In Immuno-electrophoresis and Radial Immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal Human plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified alpha-2 ceruloplasmin. No reaction is obtained with any other plasma protein component or serum. Cross-reactivity: The antiserum does not cross react with any other component of human plasma. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.
Formulation:	State: Serum State: Lyophilized, Delipidated, heat inactivated, stable whole antiserum Stabilizer: None Preservative: None



[View online »](#)

Reconstitution Method:	Restore by adding 1 ml sterile distilled water.
Concentration:	Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal Rabbit serum. No foreign proteins added.
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	ceruloplasmin (ferroxidase)
Database Link:	Entrez Gene 1356 Human P00450
Background:	Alpha-2 ceruloplasmin is copper containing alpha-2 glycoprotein with a molecular weight of 160,000. Its mean concentration in plasma of adults is 32 mg/100ml. Elevated levels are observed in pregnant women, in patients with infectious diseases and the nephritic syndrome. More than 90% of the plasma copper is bound to this protein.
Synonyms:	Ferroxidase, CP
Note:	Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other plasma proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.
Protein Families:	Druggable Genome
Protein Pathways:	Porphyryn and chlorophyll metabolism