

Product datasheet for AP21261PU-N

ZWF1 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IP, R, WB
Recommended Dilution:	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of choice. <i>Recommended Working Dilutions</i> : Non-precipitating antibody-binding techniques: 1/10,000-1/70,000.
Reactivity:	Bakers Yeast
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Glucose-6-Phosphate Dehydrogenase isolated and purified from Baker's Yeast. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	This Affinity purified antibody recognizes Glucose-6-Phosphate Dehydrogenase from Baker's
opecificity.	Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.
Formulation:	Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been
	Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. PBS, pH 7.2 stabilized with Dextran without preservatives and foreign proteins State: Aff - Purified
Formulation:	Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. PBS, pH 7.2 stabilized with Dextran without preservatives and foreign proteins State: Aff - Purified State: Lyophilized purified hyperimmune IgG fraction
Formulation: Reconstitution Method:	Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. PBS, pH 7.2 stabilized with Dextran without preservatives and foreign proteins State: Aff - Purified State: Lyophilized purified hyperimmune IgG fraction Restore by adding 0.5 ml of sterile distilled water



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	ZWF1 Rabbit Polyclonal Antibody – AP21261PU-N
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Database Link:	<u>P11412</u>
Background:	Glucose 6 Phosphate Dehydrogenase (G6PD) produces pentose sugars for nucleic acid synthesis and is the main producer of NADPH reducing power. Catalytic activity: D glucose 6 phosphate + NADP(+) → D glucono 1,5 lactone 6 phosphate + NADPH. Defects in G6PD are the cause of chronic non spherocytic haemolytic anemia (CNSHA). G6PD deficiency is the most common human enzyme deficiency; one benefit of having G6PD deficiency is that it confers a resistance to malaria.
Synonyms:	Glucose-6-phosphate 1-dehydrogenase, Glucose-6-P-Dehydrogenase

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US