

## Product datasheet for **AP21341PU-N**

### Luciferase Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IP, R, WB
Recommended Dilution:	This product is intended for use in precipitating and non-precipitating antibody-binding assays such as <i>e.g.</i> , ELISA and Western blotting and Immunofluorescence or Histochemical techniques (1/10,000-1/100,000).
Reactivity:	Photinus pyralis
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Luciferase isolated and purified from Photinus pyralis. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immuno-electrophoresis, Cross-Immuno-electrophoresis, single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.
Formulation:	PBS, pH 7.2 stabilized with Dextran without preservatives and foreign proteins. State: Aff - Purified State: Lyophilized purified IgG fraction
Reconstitution Method:	Restore by adding 0.5 ml of sterile distilled water.
Concentration:	lot specific
Purification:	Solid Phase Affinity Chromatography
Conjugation:	Unconjugated
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. If a slight precipitation occurs upon storage, this should be removed by centrifugation. Avoid repeated freezing and thawing.



[View online »](#)

**Stability:** Shelf life: one year from despatch.

**Database Link:** [P08659](#)

**Background:** Luciferase from the firefly has become one of the more widely used reporter proteins for the study of gene expression. Luciferase catalyzes a bioluminescent reaction which requires the substrate luciferin as well as  $Mg^{2+}$  and ATP. Mixing these reagents with the cell extract containing luciferase, results in a flash of light that decays rapidly. This light can be detected by a luminometer. The total light emission is proportional to the luciferase activity of the sample.