

Product datasheet for **AP21367AF-N**

phoA Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IP, R, WB
Recommended Dilution:	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques) to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of choice. <u>Working Dilutions:</u> Non-precipitating antibody-binding techniques: 1/1,000-1/80,000.
Reactivity:	Escherichia coli
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Alkaline Phosphatase isolated and purified from <i>Escherichia coli</i> . Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	Alkaline Phosphatase from <i>Escherichia coli</i> . The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.
Formulation:	PBS, pH 7.2 without preservatives and foreign proteins State: Azide Free State: Lyophilized hyperimmune IgG fraction
Reconstitution Method:	Restore by adding 1.0 ml of sterile distilled water
Concentration:	lot specific
Purification:	Ammonium Sulphate Precipitation and Ion Exchange Chromatography
Conjugation:	Unconjugated



[View online »](#)

Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. If a slight precipitation occurs upon storage, this should be removed by centrifugation. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Database Link:	P00634
Background:	Alkaline phosphatase (ALP) removes phosphate groups from the 5' end of DNA and RNA, and from proteins, at high pH. Most mammals have 4 different isozymes: placental, placental like, intestinal and non tissue specific (found in liver, kidney and bone). Tissues with particularly high concentrations of ALP include the liver, bile ducts, placenta, and bone. Damaged or diseased tissue releases enzymes into the blood, so serum ALP measurements can be abnormal in many conditions, including bone disease and liver disease.
Synonyms:	phoA, b0383, JW0374