

## Product datasheet for **AP21367BT-N**

### phoA Rabbit Polyclonal Antibody

#### Product data:

<b>Product Type:</b>	Primary Antibodies
<b>Applications:</b>	ELISA, ID, IF, IP, R, WB
<b>Recommended Dilution:</b>	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques). <i>Recommended Working Dilutions:</i> Non-precipitating antibody-binding techniques: 1/1,000-1/80,000.
<b>Reactivity:</b>	Escherichia coli
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Clonality:</b>	Polyclonal
<b>Immunogen:</b>	Alkaline Phosphatase isolated and purified from Escherichia coli. Freund's complete adjuvant is used in the first step of the immunization procedure.
<b>Specificity:</b>	Alkaline Phosphatase from <i>Escherichia coli</i> . The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.
<b>Formulation:</b>	PBS, pH 7.2 without preservatives and foreign proteins Label: Biotin State: Lyophilized purified Hyperimmune IgG fraction Molar ratio: Biotin/IgG ~ 6.6
<b>Reconstitution Method:</b>	Restore by adding 1.0 ml of sterile distilled water
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Ammonium Sulphate Precipitation and Ion Exchange Chromatography
<b>Conjugation:</b>	Biotin



[View online »](#)

<b>Storage:</b>	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
<b>Stability:</b>	Shelf life: one year from despatch.
<b>Database Link:</b>	<a href="#">P00634</a>
<b>Background:</b>	Alkaline phosphatase (ALP) removes phosphate groups from the 5' end of DNA and RNA, and from proteins, at high pH. Most mammals have 4 different isozymes: placental, placental like, intestinal and non tissue specific (found in liver, kidney and bone). Tissues with particularly high concentrations of ALP include the liver, bile ducts, placenta, and bone. Damaged or diseased tissue releases enzymes into the blood, so serum ALP measurements can be abnormal in many conditions, including bone disease and liver disease.
<b>Synonyms:</b>	phoA, b0383, JW0374