

Product datasheet for **AP21373BT-N**

Pla2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IP, R, WB
Recommended Dilution:	This antibody can be used: In precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques). <u>Recommended Dilutions:</u> non-precipitating antibody-binding assays: 1/1,000-1/40,000.
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Phospholipase A2 isolated and purified from bee venom. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	Phospholipase A2 from bee venom. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), Block Titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized Hyperimmune IgG fraction. Molar ratio: Biotin/IgG ~6.1
Reconstitution Method:	Restore by adding 1 ml of sterile distilled water.
Concentration:	lot specific
Purification:	Ammonium Sulphate Precipitation and Ion Exchange Chromatography.
Conjugation:	Biotin



[View online »](#)

Storage:	Store lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. Avoid Repeated thawing and freezing.
Stability:	Shelf life: one year from despatch.
Database Link:	P00630
Background:	<p>Phospholipase A2 catalyzes the hydrolysis of the sn-2 position of membrane glycerophospholipids to liberate arachidonic acid, a precursor of eicosanoids including prostaglandins and leukotrienes.</p> <p>This enzyme has been proposed to hydrolyze phosphatidylcholine in lipoproteins to liberate lyso-PC and free fatty acids in the arterial wall, thereby facilitating the accumulation of bioactive lipids and modified lipoproteins in atherosclerotic foci.</p> <p>Phospholipase A2 IIA can exert beneficial action in the context of infectious diseases since recent studies have shown that this enzyme exhibits potent bactericidal effects. Induction of the synthesis of Phospholipase A2 IIA is generally initiated by endotoxin and a limited number of cytokines via paracrine and/or autocrine processes.</p>
Synonyms:	Allergen Api m I, Phosphatidylcholine 2-acylhydrolase