

Product datasheet for **AP21424PU-N**

UGDH Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IP, R, WB
Recommended Dilution:	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of choice. <u>Working Dilutions:</u> Non-precipitating antibody-binding techniques: 1/100-1/9,000.
Reactivity:	Bovine
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Uridine-5'-Diphosphoglucose Dehydrogenase isolated and purified from Bovine liver. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.
Formulation:	PBS (pH 7.2) stabilized with Dextran, without preservatives and foreign proteins State: Aff - Purified State: Lyophilized purified hyperimmune IgG fraction
Reconstitution Method:	Restore by adding 0.5 ml of sterile distilled water
Concentration:	lot specific
Purification:	Solid Phase Affinity chromatography
Conjugation:	Unconjugated



[View online »](#)

Storage:	Store the antibody lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation.
Stability:	Shelf life: one year from despatch.
Database Link:	Entrez Gene 281564 Bovine P12378
Background:	UGDH is a member of the UDP-glucose/GDP-mannose dehydrogenase family and is a ubiquitously expressed protein most abundant in the liver. This protein converts UDP-glucose to UDP-glucuronate and thereby participates in the biosynthesis of glycosaminoglycans such as hyaluronan, chondroitin sulfate, and heparan sulfate. These glycosylated compounds are common components of the extracellular matrix and likely play roles in signal transduction, cell migration, and cancer growth and metastasis.
Synonyms:	UDP-glucose 6-dehydrogenase, UDP-Glc dehydrogenase, UDP-GlcDH, UDPGDH