

Product datasheet for **AP21425BT-N**

UGP2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IP, R, WB
Recommended Dilution:	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques). <u>Working Dilutions:</u> Non-precipitating antibody-binding techniques: 1/1,000-1/7,000.
Reactivity:	Bovine
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Uridine-5'-diphosphoglucose Pyrophosphorylase isolated and purified from Bovine liver. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined.
Formulation:	PBS, pH 7.2 without preservatives and foreign proteins Label: Biotin State: Lyophilized hyperimmune IgG fraction Molar ratio: Biotin/IgG ~5.0
Reconstitution Method:	Restore by adding 1.0 ml of sterile distilled water
Concentration:	lot specific
Purification:	Ammonium Sulphate Precipitation and Ion Exchange Chromatography
Conjugation:	Biotin



[View online »](#)

Storage:	Store the antibody lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation.
Stability:	Shelf life: one year from despatch.
Database Link:	Entrez Gene 281565 Bovine Q07130
Background:	UGP2 is an important intermediary in mammalian carbohydrate interconversions. It transfers a glucose moiety from glucose-1-phosphate to MgUTP and forms UDP-glucose and MgPPi. In liver and muscle tissue, UDP-glucose is a direct precursor of glycogen; in lactating mammary gland it is converted to UDP-galactose which is then converted to lactose. The eukaryotic enzyme has no significant sequence similarity to the prokaryotic enzyme.
Synonyms:	UGP1, UDP-glucose pyrophosphorylase, UGPase, UDPGP