

Product datasheet for **AP21427FC-N**

Human IgG (F(ab)2 specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Human IgG (F(ab)2 specific) Goat Polyclonal Antibody
Applications:	ELISA, ID, IF, IHC
Recommended Dilution:	<p>Direct staining of fixed cell and tissue substrates, to demonstrate the intracellular presence of free of Ig bound subunits of both kappa and lambda type. The absence of the Fc domain in the conjugate ensures minimal interaction with the tissue components and cell surfaces other than the primary antibody activity.</p> <p>This conjugate is primarily intended for use in cell surface membrane staining procedures, to identify and quantitate Ig on B cells, especially if interference by Fc activity is expected.</p> <p><i>Recommended Dilutions: 1/20-1/80.</i></p> <p>This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</p>
Reactivity:	Human
Host:	Goat
Immunogen:	<p>Purified Fab prepared from normal IgG isolated from pooled Human serum.</p> <p>Freund's complete adjuvant is used in the first step of the immunization procedure.</p>
Isotype:	IgG
Formulation:	<p>PBS, pH 7.2 without preservatives.</p> <p>Label: FITC</p> <p>State: Lyophilized Fab fragments.</p> <p>Label: Fluorescein Isothiocyanate</p> <p>Absorption emission: 492 nm / 515 nm</p> <p>Molar ratio: Fluorochrome/IgG = ~1.8</p>
Reconstitution Method:	Restore by adding 1 ml sterile di stilled water.
Concentration:	4.7 mg/ml
Purification:	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG fraction by DEAE-chromatography.



[View online »](#)

Conjugation: FITC

Storage: Store lyophilized at 2-8°C for 6 months or at -20°C long term.
After reconstitution store the antibody undiluted at 2-8°C for one month
or (in aliquots) at -20°C long term.
Avoid repeated freezing and thawing.