

Product datasheet for **AP21429TC-N**

Human IgG (Fc specific) Sheep Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Human IgG (Fc specific) Sheep Polyclonal Antibody
Applications:	ELISA, ID, IF, IHC
Recommended Dilution:	<p>Direct staining of fixed cell and tissue substrates, to demonstrate the intracellular presence of IgG. The absence of the Fc domain in the conjugate ensures minimal interaction with the tissue components and cell surfaces other than the primary antibody activity. This conjugate is primarily intended for use in cell surface membrane staining procedures, to identify and quantitate Ig on B cells, especially if interference by Fc activity is expected.</p> <p><i>Recommended Dilutions:</i> 1/10-1/40</p> <p>This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</p>
Reactivity:	Human
Host:	Sheep
Immunogen:	<p>Purified IgG prepared from pooled Human serum.</p> <p>Freund's complete adjuvant is used in the first step of the immunization procedure.</p>
Isotype:	IgG
Formulation:	<p>PBS, pH 7.2 without preservatives.</p> <p>Label: TRITC</p> <p>State: Lyophilized F(ab')₂ fragments.</p> <p>Label: Tetramethylrhodamine Isothiocyanate</p> <p>Absorption emission: 554 nm / 573 nm</p> <p>Molar ratio: Fluorochrome/IgG = ~2.7</p>
Reconstitution Method:	Restore by adding 1 ml sterile distilled water.
Concentration:	10 mg/ml
Purification:	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG fraction by DEAE-chromatography
Conjugation:	TRITC



[View online »](#)

- Storage:** Lyophilized conjugate can be shipped at ambient temperature and may be stored at 2-8°C or at -20°C for prolonged storage.
Reconstituted can be stored (into small aliquots) at -20°C.
Avoid Repeated thawing and freezing.
This product is photosensitive and should be protected from light.
- Note:** **Fluorescent marker:** Tetramethylrhodamine isothiocyanate isomer R. It has an orange-red fluorescence. To avoid nonspecific background staining, specially synthesized and exceptionally pure crystalline isomer R has been used instead of the usual racemic mixture. Although its fluorescence efficiency is less than of FITC, TRITC conjugates have the advantage of significantly less photo bleaching. This facilitates their use in quantitative cell-counting procedures.
- Adsorption:** Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.