

Product datasheet for **AP21443AF-N**

Chicken IgM (Fc specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Chicken IgM (Fc specific) Goat Polyclonal Antibody
Applications:	ID, IP
Recommended Dilution:	Can be used as unlabelled primary or secondary reagent for indirect detection of IgM at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to prepare conjugates of the user's own choice; to prepare an insoluble immunoaffinity adsorbent or a solid phase antibody reagent by coupling to an artificial carrier and as catching antibody in non-isotopic methodology and solid phase immunochemistry. When applied in any cytochemical or histochemical staining procedure or solid phase coupling technique, the optimum concentration of the IgG preparation should be established by titration before being used. <i>Recommended Working Dilutions:</i> Histochemistry: 1/50-1/250. ELISA and comparable non-precipitating antibody-binding assays: 1/500-1/5000. <i>Antibody titre:</i> Precipitin titre 1/32 when tested against pooled normal chicken serum in agar-block immunodiffusion titration.
Reactivity:	Chicken
Host:	Goat
Immunogen:	Purified IgM isolated from Chicken serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 without preservatives and foreign proteins State: Azide Free State: Lyophilized hyperimmune purified IgG fraction
Reconstitution Method:	Restore by adding 1.0 ml of sterile distilled water
Concentration:	10.0 mg/ml
Purification:	Ammonium Sulphate Precipitation and Ion Exchange Chromatography
Conjugation:	Unconjugated



[View online »](#)

Storage: Prior to and following reconstitution store the antibody at 2-8°C for one month or at -20°C for longer.
Avoid repeated freezing and thawing.

Note: **Adsorption:** Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.