

Product datasheet for **AP21450SU-N**

Equine IgM (Fc specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Equine IgM (Fc specific) Goat Polyclonal Antibody
Applications:	ID, IP
Recommended Dilution:	Can be used in precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgM in horse serum and other body fluids or to determine its concentration. To prepare an immunoabsorbent for the purification of horse IgM from serum or plasma. <u>Directions for use:</u> Immunoelectrophoresis: 2 µl or equivalent against 120 µl antiserum. Double radial immunodiffusion (Ouchterlony): a rosette arrangement with 10 µl in a 3 mm diameter center well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. <u>Antibody titre:</u> Precipitin titre 1/32 when tested against pooled normal horse serum in agar-block immunodiffusion titration.
Reactivity:	Equine
Host:	Goat
Immunogen:	Highly purified normal IgM isolated from pooled horse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formulation:	State: Serum State: Lyophilized, Delipidated, Heat inactivated, Stable Whole Serum without pr
Reconstitution Method:	Restore by adding 1 ml of sterile distilled water
Concentration:	Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Conjugation:	Unconjugated
Storage:	Prior to and following reconstitution store the antibody at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.



[View online »](#)

Note: Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.