

Product datasheet for **AP21462FC-N**

Complement C5 (C5) Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IHC, IP
Recommended Dilution:	<p>The fluorescent immunoconjugate to Human C5 is used to determine the presence and pattern using Immunohistochemical staining techniques. Determinations of individual C components can be very useful in defining the exact location of a defect.</p> <p>This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</p> <p>Working Dilutions: 1/10-1/40.</p>
Reactivity:	Human
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	<p>C5 protein isolated and purified from pooled normal human serum.</p> <p>Freund's complete adjuvant is used in the first step of the immunization</p>
Specificity:	<p>The reactivity of the antiserum is restricted to C5.</p> <p>In immunoelectrophoresis and Radial Immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal human plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified C5 protein. No reaction is obtained with any other plasma protein component or serum. The antiserum does not cross react with any other component of human plasma. Inter-species crossreactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.</p>
Formulation:	<p>PBS, pH 7.2 without preservatives.</p> <p>Label: FITC</p> <p>State: Lyophilized purified Hyperimmune IgG fraction.</p> <p>Label: Fluorescein Isothiocyanate Isomer 1</p> <p>Absorption emission: 492 nm / 515 nm</p> <p>Molar radio: Fluorescein/IgG ~2.2</p>



[View online »](#)

Reconstitution Method:	Restore by adding 1 ml of sterile distilled water.
Concentration:	lot specific
Purification:	The IgG fraction is isolated and purified from the antiserum and contains the bulk of the defined antibody specificity. It is free of other serum proteins as tested by Immunoelectrophoresis and Double Radial Immunodiffusion.
Conjugation:	FITC
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one week or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing. This product is photosensitive and should be protected from light.
Stability:	Shelf life: one year from despatch.
Gene Name:	complement component 5
Database Link:	Entrez Gene 727 Human P01031
Background:	C5 component of human complement is a plasma protein consisting of two polypeptide chains, alpha, beta, bound by disulphide bonds. It has a molecular weight of 190,000. The activated protein forms with the complement factors C6-C9 the membrane attack complex.
Synonyms:	CPAMD4, Complement component 5
Note:	Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other dog serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.
Protein Families:	Druggable Genome
Protein Pathways:	Complement and coagulation cascades, Prion diseases, Systemic lupus erythematosus