

Product datasheet for **AP21469HR-N**

Lactoferrin (LTF) Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IHC, IP, WB
Recommended Dilution:	Can be used in enzyme-immunocytochemical and immunohistochemical techniques for the detection of lactoferrin at the cellular and subcellular level in appropriately treated cell and tissue substrates; as detection reagent in nonisotopic methodology and solid phase immunochemistry (e.g. ELISA, Western blotting). This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <u>Working dilutions:</u> Histochemical and Cytochemical: 1/100 - 1/500. ELISA and comparable non-precipitating antibody-binding assays: 1/1000 - 1/7000.
Reactivity:	Human
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen has been isolated from human milk. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	In immunoelectrophoresis against human milk a single precipitin line is obtained. The antiserum does not react with any other protein component of human serum or plasma. <u>Cross-reactivity:</u> The antiserum does not cross-react with any other human plasma proteins as tested in gel-diffusion techniques. Inter-species cross-reactivity is a normal feature of antibodies to plasma or milk proteins, since homologous proteins of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail, however in double radial immunodiffusion a reaction with Rhesus monkey milk has been observed.
Formulation:	PBS, pH 7.2 without preservatives Label: HRP State: Lyophilized purified IgG fraction Label: Horseradish Peroxidase Molar ratio: Peroxidase/IgG ~1.7



[View online »](#)

Reconstitution Method:	Restore by adding 1,0 ml of sterile distilled water
Concentration:	lot specific
Purification:	Hyperimmune antisera with strong precipitating activity are selected for Fractionation by Salt-Precipitation and DEAE-Chromatography
Conjugation:	HRP
Storage:	Prior to and following reconstitution store the antibody at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	lactotransferrin
Database Link:	Entrez Gene 4057 Human P02788
Background:	Exocrine organs produce various secretions, each with its characteristic function. Proteins found in secretions may be divided into two groups: those specific for the particular secretion, and plasma proteins independent of the type of exocrine cells. Lactoferrin belongs to the first group. It is an iron containing protein with a molecular weight of 75,000 and it is antigenically different from transferrin. Lactoferrin has a slight anti-microbial action. Originally identified in milk, its presence has also been demonstrated in other secretions as saliva, semen and tears.
Synonyms:	LTF, LF, Lactoferrin, EC=3.4.21, Talalactoferrin
Note:	<u>Adsorption:</u> Immunoaffinity adsorbed using insolubilized fractions of human serum and lactoferrin-depleted human milk as required, to eliminate antibodies reacting with other human serum or milk proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.
Protein Families:	Druggable Genome, Protease, Secreted Protein