

Product datasheet for **AP21471AF-N**

Fibrinogen Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IF, IP, WB
Recommended Dilution:	Can be used: As unlabelled primary or secondary antibody reagent for the indirect detection of Fibrinogen in Rat cells, tissues and body fluids in Immunofluorescence and Immunoenzyme methods. For production of immunoconjugates with a selected marker. To prepare insoluble immunoaffinity adsorbents by coupling to an artificial carrier. As catching or detection reagent in non-isotopic methodology and solid phase immunochemistry (e.g. ELISA, Western blotting). <u>Recommended Dilutions:</u> Histochemistry: 1/50-1/250. ELISA and comparable non-precipitating antibody-binding assays: 1/500-1/5,000.
Reactivity:	Rat
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fibrinogen is isolated from fresh plasma after removing Prothrombin. Freund's complete adjuvant is used in the first step of the immunization procedure.
Specificity:	The reactivity of the antiserum is restricted to Fibrinogen. In Immunoelectrophoresis and Radial immunodiffusion (Ouchterlony), using various antiserum concentrations against normal rat plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified fibrinogen. No reaction is obtained with any other plasma protein component or serum. However, the antiserum may also react with fibrin monomers, circulating fibrinopeptides and fibrin degradation products. The antiserum does not cross-react with any other component of rat plasma. Inter-species crossreactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.
Formulation:	PBS, pH 7.2 without preservatives. State: Azide Free State: Lyophilized purified IgG fraction.



[View online »](#)

Reconstitution Method:	Restore by adding 1 ml of sterile distilled water.
Concentration:	lot specific
Purification:	The IgG fraction is isolated and purified from the antiserum and contains the bulk of the defined antibody specificity. It is free of other serum proteins as tested by Immunoelectrophoresis and double Radial Immunodiffusion.
Conjugation:	Unconjugated
Storage:	Store lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. Avoid Repeated thawing and freezing.
Stability:	Shelf life: one year from despatch.
Background:	Fibrinogen (clotting factor I) is a heat labile beta glycoprotein present in plasma. It is the precursor of fibrin, which is the key protein constituting the network of the blood clot. Thrombin converts fibrinogen to fibrin by limited proteolysis. Fibrin monomers polymerize to fibrin which is stabilized by cross-linking.
Synonyms:	FGA, FGB, FGG
Note:	Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other with other plasma proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.