

Product datasheet for **AP21471FC-N**

Fibrinogen Goat Polyclonal Antibody

Product data:

| | |
|-----------------------|--|
| Product Type: | Primary Antibodies |
| Applications: | ELISA, ID, IF, IHC |
| Recommended Dilution: | Suitable for the direct detection of Fibrinogen in Rat cells, tissues and body fluids in Immunofluorescence. Also suitable as detection reagent in non-isotopic methodology and Solid Phase Immunochemistry (e.g. ELISA). This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <i>Recommended Dilutions: 1/10-1/40.</i> |
| Reactivity: | Rat |
| Host: | Goat |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Fibrinogen is isolated from fresh plasma after removing Prothrombin. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Specificity: | The reactivity of the antiserum is restricted to Fibrinogen. In Immunoelectrophoresis and Radial Immunodiffusion (Ouchterlony), using various antiserum concentrations against normal Rat plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified Fibrinogen. No reaction is obtained with any other plasma protein component or serum. However, the antiserum may also react with Fibrin monomers, circulating fibrinopeptides and Fibrin degradation products. Cross-Reactivity: The antiserum does not cross-react with any other component of Rat plasma. Inter-species crossreactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail. |



[View online »](#)

| | |
|-------------------------------|--|
| Formulation: | PBS, pH 7.2 without preservatives and foreign proteins Label: FITC State: Lyophilized purified IgG fraction Label: Fluorescein Isothiocyanate Isomer 1 Absorption emission: 492 nm / 515 nm Molar ratio: Fluorochrome/IgG ~2.2 |
| Reconstitution Method: | Restore by adding 1 ml of sterile distilled water |
| Concentration: | lot specific |
| Purification: | Hyperimmune antisera with strong precipitating activity are selected for Fractionation by Salt Precipitation and purification of the IgG fraction by DEAE-Chromatography |
| Conjugation: | FITC |
| Storage: | Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one week or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| Background: | Fibrinogen (clotting factor I) is a heat labile beta glycoprotein present in plasma. It is the precursor of fibrin, which is the key protein constituting the network of the blood clot. Thrombin converts fibrinogen to fibrin by limited proteolysis. Fibrin monomers polymerize to fibrin which is stabilized by cross-linking. |
| Synonyms: | FGA, FGB, FGG |
| Note: | Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other with other plasma proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum. |