

Product datasheet for **AP21481AF-N**

Mouse IgD (Fc specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Mouse IgD (Fc specific) Goat Polyclonal Antibody
Applications:	ELISA, ID, IF, IP, WB
Recommended Dilution:	Can be used as unlabelled primary or secondary reagent for indirect detection techniques, to prepare conjugates with markers of the user's own choice, to prepare an insoluble Immunoaffinity adsorbent or a solid phase antibody reagent by coupling to an artificial carrier and as catching or detection antibody in non-isotopic methodology and solid phase Immunochemistry. When applied in any Cytochemical or Histochemical procedure or solids phase coupling technique, the optimum concentration of the IgG preparation should always be established by titration. <i>Recommended Working Dilutions:</i> Histochemical Use: 1/50-1/250. ELISA and comparable non-precipitating antibody-binding assays: 1/500-1/5000.
Reactivity:	Mouse
Host:	Goat
Immunogen:	Purified polyclonal and monoclonal IgD isolated from BALB/C and C57BL Mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 without preservatives State: Azide Free State: Lyophilized purified Hyperimmune IgG fraction
Reconstitution Method:	Restore by adding 1.0 ml sterile distilled water
Concentration:	10.0 mg/ml
Purification:	DEAE Chromatography
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.



[View online »](#)

Note: ***Adsorption:*** Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.