

Product datasheet for **AP21481TC-N**

Mouse IgD (Fc specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Mouse IgD (Fc specific) Goat Polyclonal Antibody
Applications:	ELISA, ID, IF, IHC, IP
Recommended Dilution:	In immunocytochemical and immunohistochemical staining for the detection of IgD at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to identify and measure IgD in mouse serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <u>Recommended working dilutions:</u> 1/10 - 1/40, depending on the method used.
Reactivity:	Mouse
Host:	Goat
Immunogen:	Pools of purified homogenous IgD isolated from BALB/C and C57Bl mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 without preservatives or foreign proteins Label: TRITC State: Lyophilized purified Ig fraction Label: Tetramethylrhodamine Isothiocyanate Absorption emission: 554 nm / 573 nm Molar ratio: Fluorochrome/IgG ~1.7
Reconstitution Method:	Restore by adding 1,0 ml sterile distilled water
Concentration:	10,0 mg/ml
Purification:	Hyperimmune antisera with strong precipitating activity are selected for fractionation and purification of the IgG (7S) fraction containing the bulk of the defined antibody specificity. It is free of other serum proteins as tested by immunoelectrophoresis.
Conjugation:	TRITC



[View online »](#)

- Storage:** Prior to and following reconstitution store the antibody at 2-8°C for one month or at -20°C for longer.
Avoid repeated freezing and thawing.
- Note:** Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.