

Product datasheet for **AP21485HR-N**

Mouse IgE (Fc specific) Rabbit Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Mouse IgE (Fc specific) Rabbit Polyclonal Antibody
Applications:	ELISA, ID, IF, IHC, IP, WB
Recommended Dilution:	Can be used in enzyme-immunocytochemical and immunohistochemical staining for the detection of IgE at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgE antibodies in mouse serum or other body fluids; in non-isotopic assay methodology (e.g. ELISA) to identify and measure IgE in mouse serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <u>Recommended working dilutions:</u> Histochemical and Cytochemical: 1/100 - 1/500. ELISA and comparable non-precipitating antibody-binding assays: 1/500 - 1/5000.
Reactivity:	Mouse
Host:	Rabbit
Immunogen:	Purified homogenous IgE isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 without preservatives and foreign proteins Label: HRP State: Lyophilized purified IgG fraction Label: Horseradish Peroxidase <u>Enzyme marker:</u> Horseradish peroxidase enriched for isoenzyme C (RZ=3.2) <u>Conjugation procedure:</u> Conjugation is carried out using a proprietary modification of the periodate technique for the binding to peroxidase, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilized to assure stability and long shelf life Molar ratio: Peroxidase/IgG: ~1.7
Reconstitution Method:	Restore by adding 0.5 ml sterile distilled water.
Concentration:	10,0 mg/ml



[View online »](#)

Purification:	Immunoaffinity Chromatography
Conjugation:	HRP
Storage:	Prior to and following reconstitution store the antibody at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.
Note:	<u>Adsorption:</u> Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.