

Product datasheet for **AP21493BT-N**

Rat IgG2ab (subclass specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Rat IgG2ab (subclass specific) Goat Polyclonal Antibody
Applications:	ELISA, ID, IF, IHC, WB
Recommended Dilution:	<p>Can be used:</p> <p>In immunocytochemical and Immunohistochemical staining for the detection of IgG2 at the cellular and subcellular level of appropriately treated cell and tissue substrates.</p> <p>To demonstrate circulating IgG2 antibodies in serodiagnostic microbiology and autoimmune diseases.</p> <p>To identify a specific antigen using a reference antibody of Rat origin known to be of the IgG2 isotype in the middle layer of the indirect test procedure.</p> <p>In non-isotopic assay methodology (e.g. ELISA) to measure IgG2 in rat serum or other body fluids.</p> <p>As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</p> <p><i>Recommended Working Dilutions:</i></p> <p>Histochemical and Cytochemical Use: 1/100-1/250.</p> <p>ELISA and comparable non-precipitating antibody-binding assays between: 1/2.000-1/10,000.</p>
Reactivity:	Rat
Host:	Goat
Immunogen:	<p>Pools of highly purified homogenous IgG2a and IgG2b isolated from rat serum.</p> <p>Freund's complete adjuvant is used in the first step of the immunization procedure.</p>
Isotype:	IgG
Formulation:	<p>PBS, pH 7.2</p> <p>Label: Biotin</p> <p>State: Lyophilized purified IgG fraction</p> <p>Preservative: None, as it may interfere with the antibody activity</p> <p>Molar ratio: Biotin/IgG: ~6.4</p>
Reconstitution Method:	Restore by adding 1.0 ml sterile distilled water.
Concentration:	10.0 mg/ml



[View online »](#)

Purification:	Hyperimmune antisera with strong precipitating activity are selected for fractionation and purification of the IgG fraction containing the bulk of the defined antibody specificity. It is free of other serum proteins as tested by immunoelectrophoresis.
Conjugation:	Biotin
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Note:	Marker: N-Hydroxysuccinimidobiotin. Conjugation procedure: A proprietary technique for the binding of biotin, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilized to assure stability and long shelf life. Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.