

Product datasheet for AP22860PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

Phone: +1-888-26/-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase 1 (CASP1) (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IHC, WB

Recommended Dilution: ELISA.

Immunohistochemistry on Paraffin Sections: 5 µg/ml.

Western Blot: 0.5 µg/ml.

Reactivity: Human
Host: Rabbit

Clonality: Polyclonal

Immunogen: Recombinant rat HO-1 (Hsp32) lacking the membrane spanning region

Specificity: This antibody reacts to Caspase 1 (CASP1).

Formulation: PBS containing 0.02% sodium azide.

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Immunoaffinity Chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Gene Name: caspase 1

Database Link: Entrez Gene 834 Human

P29466

Background: Caspase 1, also called interleukin-1 beta converting enzyme, is a cysteine C14 endopeptidase

able to process pro-interleukin-1 beta into its active form. Caspase 1 is identical to Caenorhabditis elegans cell death gene ced-3 and may be involved in apoptotic signalling.

Inhibition of caspase-1 in mice models extended survival and delayed appearance of

neuronal inclusions and symptoms of Huntingdon disease.

Synonyms: CASP-1, CASP1, IL1BC, IL1BCE, ICE, p45

Protein Families: Druggable Genome, Protease







Protein Pathways:

Amyotrophic lateral sclerosis (ALS), Cytosolic DNA-sensing pathway, NOD-like receptor signaling pathway


Product images:



Western blot analysis of caspase-1 in human heart cell lysate with caspase-1 antibody (AP22860PU-N) at (A) 0.5, (B) 1, and (C) 2 ug/ml.



Human Heart (formalin-fixed, paraffinembedded) stained with CASP1 antibody AP22860PU-N at 5 ug/ml followed by biotinylated goat anti-rabbit IgG secondary antibody, alkaline phosphatase-streptavidin and chromogen.



Human Testis (formalin-fixed, paraffinembedded) stained with CASP1 antibody AP22860PU-N at 5 ug/ml followed by biotinylated goat anti-rabbit IgG secondary antibody, alkaline phosphatase-streptavidin and chromogen.