

#### Product datasheet for AP23297PU-N

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **Angpt2 (C-term) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

**Recommended Dilution:** Western blot: At 1-2µg/ml with the appropriate system to detect ANG2 in cells and tissues.

Immunohistochemistry on Paraffin Sections: At 1-2µg/ml to detect ANG2 in formalin fixed

and paraffin embedded.

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a sequence mapping near the C-terminal (454-468) of

Human Angiopoietin-2.

**Specificity:** This antibody detects Angiopoietin-2 near C-term.

No cross reactivity with other proteins.

**Formulation:** 0.9mg NaCl, 0.2mg Na2HPO4

State: Aff - Purified

State: Lyophilized purified Ig fraction

Stabilizer: 5mg BSA

Preservative: 0.05mg Thimerosal, 0.05mg Sodium Azide

**Reconstitution Method:** 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

**Purification:** Immunoaffinity Chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** Mus musculus angiopoietin 2 (Angpt2)

Database Link: Entrez Gene 285 HumanEntrez Gene 89805 RatEntrez Gene 11601 Mouse

O35608






#### Background:

Angiopoietin 1 and Angiopoietin 2 are important for development of the endothelium, by regulating tyrosine phosphorylation of the membrane receptor Tie 2. Angiopoietin 2 is only 60% homologous with Angiopoietin 1. Angiopoietin-2 is a naturally occurring antagonist of angiopoietin-1 that competes for binding to the TIE2 receptor and blocks ANGPT1-induced TIE2 autophosphorylation. Angiopoietin 1 binding to Tie 2 causes phosphorylation of the receptor. Angiopoietin 2 competes for this binding, and thus blocks receptor phosphorylation. Angiopoietin 2 expression occurs at sites of vascular remodelling: dorsal aorta and major aortic branches, ovary, placenta and uterus.


Synonyms:

ANG-2, ANG2, ANGPT2

## **Product images:**



Western blot analysis of rat brain tissue lysis using ANG2 antibody



Immunohistochemical analysis of paraffin embedded Human placencta sections using ANG2 antibody