

Product datasheet for AP23335PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

OPCML (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: At 1µg/ml with the appropriate system to detect OPCML in cells and tissues.

Immunohistochemistry on poraffin sections: At 1-2µg/ml to detect OPCML in formalin fixed

and paraffin embedded tissues.

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide mapping at the C-terminal of human OPCML

Specificity: This antibody detects OPCML / OBCAM (C-term).

Formulation: 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3

State: Aff - Purified

State: Lyophilized Ig fraction

Reconstitution Method: 0.2ml of distilled water will yield a concentration of 500µg/ml.

Purification: Immunogen affinity purified

Conjugation: Unconjugated

Storage: Store at 2 - 8 °C for up to one month or (in aliquots) at -20 °C for longer. Avoid repeated

freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: opioid binding protein/cell adhesion molecule-like

Database Link: Entrez Gene 116597 RatEntrez Gene 330908 MouseEntrez Gene 4978 Human

Q14982






Background:

Opioid-binding protein/cell adhesion molecule-like (OPCML), (also called opioid-binding cell adhesion molecule (OBCAM)). OPCML is a member of the IgLON family of immunoglobulin (Ig)domain-containing glycosylphosphatidylinositol (GPI)-anchored cell adhesion molecules, as an excellent candidate for the 11q25 ovarian cancer TSG in EOC. The OPCML gene comprises 7 exons, spans approximately 600 kb, and is transcribed from telomere to centromere . And due to the lack of transmembrane domains necessary for signal transduction, it is improbable that OBCAM acts independently as an opioid receptor; more likely, it plays an important accessory role in opioid receptor function.


Synonyms: Opioid-binding cell adhesion molecule

Protein Families: Druggable Genome, Transmembrane

Product images:



OPCML Polyclonal Antibody



OPCML Polyclonal Antibody