

Product datasheet for **AP23431PU-N**

Visfatin (NAMPT) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	Sandwich ELISA: To detect Human Visfatin by Sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5-2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human Visfatin (AP23431BT-N or AP23431BT-S) as a detection antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant Human Visfatin. Western Blot: To detect Human Visfatin by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Human Visfatin is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions. Immunohistochemistry: The antibody stained FFPE Sections of human ovary malignant adeocarcinoma. An HRP-labelled polymer detection system was used with a DAB chromogen). Heat induced antigen retrieval with pH 6.0 Sodium Citrate buffer is recommended. <i>Working Dilution:</i> 7.5-15 ng/ml, overnight incubation at 4°C.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure E.coli derived recombinant Human Visfatin.
Specificity:	Recognizes Visfatin
Formulation:	PBS, pH 7.2 without preservatives State: Aff - Purified State: Lyophilized (sterile filtered) purified Ig fraction
Reconstitution Method:	Centrifuge vial prior to opening. Restore in sterile water to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity Chromatography employing an immobilized Human Visfatin matrix.
Conjugation:	Unconjugated



[View online »](#)

Storage:	Store the lyophilized antibody at -20°C. Following reconstitution it is stable for two weeks at 2-8°C. Frozen aliquots are stable for 6 months when stored at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	nicotinamide phosphoribosyltransferase
Database Link:	Entrez Gene 10135 Human P43490
Background:	<p>Visfatin, a newly identified adipocytokine, which is predominantly secreted from visceral adipose tissue both in humans and mice. Visfatin corresponds to a protein identified previously as pre-B cell colony-enhancing factor (PBEF). Visfatin exerted insulin-mimetic effects in cultured cells, various insulin-sensitive tissues such as liver, muscle, and fat, and lowered plasma glucose levels in mice. Mice heterozygous for a targeted mutation in the visfatin gene had modestly higher levels of plasma glucose relative to wild-type littermates. visfatin binds to and activates the insulin receptor. Which may lead to new insights into glucose homeostasis and/or new therapies for metabolic disorders such as diabetes. Excess adiposity is the most important risk in the development of insulin resistance and type 2 diabetes mellitus (T2DM). Adipose tissue produces several proteins (adipocytokines) such as leptin, adiponectin, resistin, tumor necrosis factor-alpha, and IL-6, that modulate insulin sensitivity and appear to play an important role in the pathogenesis of insulin resistance, diabetes, dyslipidemia, inflammation, and atherosclerosis. However, the mechanisms by which fat tissue induces insulin resistance and the role of adipocytokines in the pathogenesis of T2DM have not been well established. Visfatin, also known as pre-B cell colony-enhancing factor (PBEF), is a cytokine that is highly expressed in visceral fat and was originally isolated as a secreted factor that synergizes with IL-7 and stem cell factors to promote the growth of B cell precursors. Visfatin homologs have been identified in carp, invertebrate mollusks, and bacteria, as well as in vertebrates, including humans and the mouse. It has been postulated to play a role in innate immunity.</p> <p>Visfatin exerts insulin-mimetic effects that are dose-dependent and quantitatively similar to those of insulin in stimulating muscle and adipocyte glucose transport, and in inhibiting hepatocyte glucose production. Intravenous injection of recombinant visfatin in mice decreased plasma glucose in a dose-dependent fashion. In keeping with its insulin-mimetic effects, visfatin was as effective as insulin in reducing hyperglycemia in insulin-deficient diabetic mice. Visfatin was also found to be bound to and activate insulin receptor, causing receptor phosphorylation and the activation of downstream signaling molecules. However, visfatin and insulin did not compete for binding to the insulin receptor, indicating that the two proteins were recognized by different regions of the receptor. Thus, visfatin might play a role in glucose homeostasis and dysregulation in biosynthesis or signal transduction, and might contribute to the pathogenesis of diabetes.</p>
Synonyms:	NAmPRTase, PBEF, PBEF1
Protein Families:	Druggable Genome

Protein Pathways: Nicotinate and nicotinamide metabolism