

## Product datasheet for **AP23432BT-S**

### CCL28 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	<b>Direct ELISA:</b> To detect Human MEC by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25–1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2–0.4 ng/well of recombinant Human MEC. <b>Sandwich ELISA:</b> To detect Human MEC by Sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.25-1.0 µg/ml of this antibody is required. This Biotin conjugated, in conjunction with Purified Anti-Human Visfatin (AP23432PU-N or AP23432PU-S) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant Human MEC. <b>Western Blot:</b> To detect Human MEC by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Human MEC is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure recombinant Human MEC
Specificity:	Recognizes MEC.
Formulation:	PBS, pH 7.2 without preservatives State: Aff - Purified State: Lyophilized (sterile filtered) purified Ig fraction
Reconstitution Method:	Centrifuge vial prior to opening. Restore in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity Chromatography
Conjugation:	Unconjugated



[View online »](#)

<b>Storage:</b>	Store the lyophilized antibody at -20°C. Following reconstitution it is stable for two weeks at 2-8°C. Frozen aliquots are stable for 6 months when stored at -20°C. Avoid repeated freezing and thawing.
<b>Stability:</b>	Shelf life: One year from despatch.
<b>Gene Name:</b>	C-C motif chemokine ligand 28
<b>Database Link:</b>	<a href="#">Entrez Gene 56477 Human</a> <a href="#">Q9NRJ3</a>
<b>Background:</b>	CCL28, also known as chemokine (C-C motif) ligand 28 or mucosae-associated epithelial chemokine (MEC), belongs to the subfamily of small cytokine CC proteins. CCL28 is expressed by columnar epithelial cells in the gut, lung, breast and the salivary glands and drives the mucosal homing of T and B lymphocytes that express CCR10, and the migration of eosinophils expressing CCR3. This chemokine is constitutively expressed in the colon, but its levels can be increased by pro-inflammatory cytokines and certain bacterial products implying a role in effector cell recruitment to sites of epithelial injury.
<b>Synonyms:</b>	C-C motif chemokine 28, SCYA28
<b>Protein Families:</b>	Druggable Genome, Secreted Protein
<b>Protein Pathways:</b>	Chemokine signaling pathway, Cytokine-cytokine receptor interaction