

Product datasheet for AP23440SU-S

Uroguanylin (GUCA2B) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


| Product Type: | Primary Antibodies |
|------------------------|--|
| Applications: | IHC, WB |
| Recommended Dilution: | Western blot: 1/20,000 (Ref.1). Immunohistochemistry on Frozen Sections: 1/2,000 (Ref.1). |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Immunogen: | Synthetic Human Uroguanylin |
| Specificity: | This antibody recognizes Synthetic Human Uroguanylin. |
| Formulation: | State: Serum State: Lyophilized Serum |
| Reconstitution Method: | Restore in aqua bidest to initial volume. |
| Conjugation: | Unconjugated |
| Storage: | Store lyophilized at 2-8°C and reconstituted at -20°C. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: One year from despatch. |
| Gene Name: | guanylate cyclase activator 2B |
| Database Link: | <u>Entrez Gene 2981 Human</u> <u>Q16661</u> |
| Synonyms: | UGN |
| Protein Families: | Druggable Genome, Secreted Protein, Transmembrane |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Western nlot analysis of AP23440SU specificity. Prostatic fluid from healthy donors was separated by SDS-PAGE und reducing conditions and immunoblotted with AP23440SU (1/20,000). Lane 1: AP23440SU detects Uroguanylin as a band migrating at 12-13 kDa. Lane 2: Negative Control incubated with non-immune Serum. Arrow: Aprotinin, 9 kDa. Double Arrow: Lysozyme, 16.8 kDa. Maake C et al. (2003). Histochem Cell Biol 119 (1):69-76.



Immunohistochemistry of Uroguanylin staining in Cyryosection of Human Prostate. The section was Fixed in Acetone, incubated with AP23440SU (1/2000) 72h at 4°C and detected using Biotin-Streptavidin technique. 3, 3'-di-aminobenzidine was used as the Chromogen. This antibody detects Uroguanylin in luminal part of grandular epithelial cells and in cell protrusions (arrows). Maake C et al. (2003). Histochem Cell Biol 119 (1):69-76.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US