

Product datasheet for **AP26075PU-S**

PKM2 (PKM) (Isoform M1) Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	Western blot: 1/1000. Immunofluorescence.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide sequence around aa. 399~403 derived from Human PKM1.
Specificity:	This antibody detects endogenous levels of total PKM1 protein.
Formulation:	Phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% Sodium Azide and 50% Glycerol State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	lot specific
Purification:	Affinity-chromatography using epitope-specific peptide
Conjugation:	Unconjugated
Storage:	Upon receipt, store undiluted (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	60 kDa
Gene Name:	pyruvate kinase, muscle
Database Link:	Entrez Gene 18746 Mouse Entrez Gene 5315 Human P14618




[View online »](#)

Background:	Glycolytic enzyme that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP. Stimulates POU5F1-mediated transcriptional activation. Plays a general role in caspase independent cell death of tumor cells. The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production. The transition between the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and survival.
Synonyms:	PK2, PK3, PKM, CTHBP, M2-PK, THBP1, OIP3, OIP-3, Pyruvate kinase 2/3, Pyruvate kinase M1/M2, Pyruvate kinase muscle
Protein Families:	Druggable Genome
Protein Pathways:	Glycolysis / Gluconeogenesis, Metabolic pathways, Purine metabolism, Pyruvate metabolism, Type II diabetes mellitus

Product images:

Western blot analysis of extracts from U87 cells using PKM1 Antibody.



Immunofluorescence staining of methanol-fixed MEF cells using PKM1 Antibody.