

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for AP26422AF-N

## NFKB1 pSer337 Rabbit Polyclonal Antibody

## **Product data:**

Product Type:	Primary Antibodies
Applications:	ELISA
Recommended Dilution:	<b>ELISA:</b> The antibody (approx. 1:100,000 - 1:400,000 diluted) can detect the serine 337 phosphorylated NF-κB p105/p50 peptide coated on ELISA plate at 100 ng/well.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	A synthetic peptide derived from Serine337 phosphorylated NF-кВ p105/p50, conjugated with KLH
Specificity:	This antibody is reactive to the synthetic NF-кВ p105/p50 peptide (QLRRKSpDLETSEP) on Mouse, human and other species with consensus serine 337 phosphorylated NF- кВ p105/p50 peptide.
Formulation:	0.01M PBS, pH 7.4 State: Azide Free State: Lyophilized Ig fraction
Reconstitution Method:	Restore with double distilled water to adjust the final concentration to 1.00 mg/ml.
Purification:	Protein G affinity
Conjugation:	Unconjugated
Storage:	Store lyophilized at 2-8°C for 6 month or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	nuclear factor kappa B subunit 1
Database Link:	<u>Entrez Gene 4790 Human</u> <u>P19838</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	NFKB1 pSer337 Rabbit Polyclonal Antibody – AP26422AF-N
Background:	NF-κB (nuclear factor kappa-light-chainenhancer of activated B cells) is a transcription factor that controls the transcription of many genes that are associated with immuno-response, antiapoptosis, neuroplasticity and memory. There are five NF-κB proteins in mammals: RelA (NF-κB-p65), RelB, c-Rel, NF-κB (NF-κB p105), and NF-κB2 (NF-κB p100). They form a variety of homodimers and heterodimers, each of which activates its own distinctive set of genes. P65 has been shown to be involved with DNA binding. The phosphorylation of p65 at Ser468 is inducible by TNF. Ser468 phosphorylation mediates p65 ubiquitination and the termination of related gene expression.
Synonyms:	NFKB1, KBF1, EBP-1, EBP1, NF-kappa-B p50

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US