

## Product datasheet for **AP31439FC-N**

### Monkey IgG (H+L chain) Goat Polyclonal Antibody

#### Product data:

Product Type:	Secondary Antibodies
Product Name:	Monkey IgG (H+L chain) Goat Polyclonal Antibody
Applications:	ELISA, IF, IHC, IP
Recommended Dilution:	To identify and measure IgG, antigen or antibody, at the cellular and subcellular level by immunofluorescence staining of appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of monkey origin in the middle layer of the indirect test procedure. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <u>Working dilutions</u> are usually between 1/20 and 1/80.
Reactivity:	Monkey
Host:	Goat
Immunogen:	Purified normal IgG isolated from pooled rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 No preservative added. - No foreign proteins added. Label: FITC State: Lyophilised hyperimmune Ig fraction Label: Fluorescein isothiocyanate isomer 1 Absorption emission: 492nm/515nm Molar radio: 1,5
Reconstitution Method:	Restore with 2 ml sterile distilled water.
Concentration:	10 mg/ml
Purification:	DEAE-column Chromatography
Conjugation:	FITC



[View online »](#)

**Storage:**

Prior to reconstitution store at 2-8°C.  
Following reconstitution store undiluted at 2-8°C for one week  
or (in aliquots) at -20°C for longer.  
Avoid repeated freezing and thawing.