

Product datasheet for **AP31457BT-N**

Pigeon IgG (H+L chain) Rabbit Polyclonal Antibody

Product data:

| | |
|-----------------------|--|
| Product Type: | Primary Antibodies |
| Applications: | ELISA, IF, IHC, WB |
| Recommended Dilution: | <p>Can be used in immunocytochemical and immunohistochemical staining to identify and measure IgG, antigen or antibody, at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of pigeon origin in the middle layer of the indirect test procedure. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</p> <p>Working dilutions fro histochemical and cytochemical use are usually between 1/100 and 1/500.</p> <p>In ELISA and comparable non-precipitating antibody-binding assays between 1/2000 and 1/15000.</p> |
| Reactivity: | Pigeon |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Purified normal IgG isolated from pooled pigeon serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |



[View online »](#)

| | |
|-------------------------------|--|
| Specificity: | <p>The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in pigeon serum, as tested by immunoelectrophoresis and double radial immunodiffusion.</p> <p><u>Cross-reactivity:</u> Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. The degree of cross-reactivity depends in addition to the phylogenetic relationship also on the concentration of the reactants and the sensitivity of the assay format. Cross-reactivity of this antiserum has not been tested in detail, however in immunoelectrophoresis a reaction with chicken, duck and turkey has been observed.</p> |
| Formulation: | <p>PBS, pH 7.2 No preservative added - No foreign proteins added Label: Biotin State: Lyophilised hyperimmune Ig fraction Molar ratio: 3,3</p> |
| Reconstitution Method: | Restore with 1 ml sterile distilled water |
| Concentration: | lot specific |
| Purification: | DEAE-column Chromatography |
| Conjugation: | Biotin |
| Storage: | <p>Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one week or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.</p> |
| Stability: | Shelf life: one year from despatch. |
| Synonyms: | Pigeon Immunoglobulin G |