

Product datasheet for AP31466SU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Duck Serum Proteins Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ID, IP, R

Recommended Dilution: In precipitating techniques as immunoelectrophoresis and radial immunodiffusion

(Ouchterlony) to identify the serum protein pattern, or the presence or absence of an individual component. To evaluate the purity of an isolated serum protein including immunoglobulins. Since immunoprecipitation depends on a correct antigen/antibody concentration ratio (zone of equivalence) in the gel medium, the protein analysis by immunoelectrophoresis of serum or any other biological fluid or protein fraction should include different proportions of the reactants. It is not possible to obtain an optimal protein pattern in a single analysis. The electroendosmosis effect of different types of agar on proteins with a different net charge can be used to optimize the resolution power of the test system. Agar contains sufficient positively charged ions to optimize the resolution of the proteins in the beta-gamma regions, while the alpha regions will become more dense. Highly purified agar with low electroendosmosis favours the resolution of the proteins in the alpha regions, while the major components in the beta-gamma region can still be identified.

Directions for use:

In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l serum samples (neat and serially diluted in

2 mm diameter peripheral wells.

Reactivity: Duck
Host: Rabbit
Clonality: Polyclonal

Immunogen: Pooled whole duck serum and partly purified serum fractions. Freund's complete adjuvant is

used in the first step of the immunization procedure.

Specificity: In immunoelectrophoresis against pooled serum and enriched serum proteins fractions

precipitation can be observed of not less than 12 individual proteins components.

<u>Cross-reactivity:</u> Inter-species cross-reactivity is a normal feature of antibodies to animal proteins since homologous proteins of different species frequently share antigenic

determinants. This antiserum has not been adsorbed for such cross-reactivity. Consequently

it is not species-specific.





Duck Serum Proteins Rabbit Polyclonal Antibody - AP31466SU-N

Formulation: No preservative added - No foreign proteins added

State: Serum

State: Delipidated, heat inactivated, lyophilized, stable whole antiserum

Reconstitution Method: Restore with 1 ml sterile distilled water.

Conjugation: Unconjugated

Storage: Prior to reconstitution store at 2-8°C.

Following reconstitution store undiluted at 2-8°C for one week

or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.