

# Product datasheet for AP31512FC-N

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Alb Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ID, IF, R

**Recommended Dilution:** Can be used as reagent for the direct detection of Albumin in Mouse cells, tissues and body

fluids in Immunofluorescence techniques.

This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the

specific signal.

Recommended Dilutions: 1/20-1/80.

Reactivity: Mouse
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Albumin is isolated from Mouse serum by Sequential Precipitation and purified by Ion

Exchange Chromatography and Affinity Chromatography.

Freund's complete adjuvant is used in the first step of the immunization procedure.

**Specificity:** The defined antibody specificity is directed to Albumin as tested against Mouse sera. In

Immunoelectrophoresis and double radial Immunodiffusion (Ouchterlony), using various antiserum concentrations against appropriate concentrations of the immunogen, a single characteristic precipitin line is obtained which shows a reaction of identity with the precipitin

lines obtained against Mouse serum and the purified Albumin.

**Cross-reactivity:** Inter-species cross-reactivity is a normal feature of antibodies to serum proteins, since homologous proteins of different species frequently share antigenic

determinants. The degree of cross-reactivity is also dependent on the concentrations of the

reactants and the sensitivity of the assay arrangement.

This antiserum has been tested for cross-reactivity in Double Radial Immunodiffusion with

the following results:

Bovine (-), Chicken (-), Canine (-), Goat (-), Guinea Pig (++), Hamster (+), Horse (+), Human (+),

Monkey (+), Rat (++), Sheep (-), Swine (+).





#### Alb Rabbit Polyclonal Antibody - AP31512FC-N

**Formulation:** PBS, pH 7.2 without preservatives and foreign proteins

Label: FITC

State: Lyophilized purified hyperimmune IgG fraction

Label: Fluorescein isothiocyanate isomer 1 Absorption emission: 492 nm / 515 nm Molar radio: Fluorescein/IgG ~1.7

**Reconstitution Method:** Restore by adding 1 ml of sterile distilled water

**Concentration:** lot specific

**Purification:** Hyperimmune with strong precipitating activity are selected for fractionation by Salt

Precipitation and the purification of the IgG fraction by DEAE Chromatography

Conjugation: FITC

**Storage:** Prior to reconstitution store at 2-8°C.

Following reconstitution store undiluted at 2-8°C for one week

or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: albumin

Database Link: Entrez Gene 11657 Mouse

P07724

**Background:** Albumin is a soluble, monomeric protein which comprises about one half of the blood serum

protein. Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Mutations in this gene on chromosome 4 result in various anomalous proteins. Albumin is a globular unglycosylated serum protein of molecular weight 65,000. The human albumin gene is 16,961 nucleotides long from the putative 'cap' site to the first poly(A) addition site. It is split into 15 exons which are symmetrically placed within the 3 domains that are thought to have arisen by triplication of a single primordial domain. Albumin is synthesized in the liver as preproalbumin which has an N terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in turn cleaved in the Golgi vesicles to

produce the secreted albumin.

Synonyms: ALB, BSA, HSA, Serum Albumin

**Note:** <u>Conjugation procedure:</u> A proprietary technique for the binding to FITC is used, followed by

several purification steps to remove free reactants and protein aggregates. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilized to

assure stability and long shelf life.