

Product datasheet for **AP31539BT-N**

Canine IgA (Fc specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Canine IgA (Fc specific) Goat Polyclonal Antibody
Applications:	ID, IP
Recommended Dilution:	Can be used in immunocytochemical and immunohistochemical staining of IgA at the cellular and subcellular level of appropriately treated cell and tissue substrates; to demonstrate circulating IgA antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of dog origin known to be of the IgA isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgA in dog serum or other body fluids. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. Antisera to IgA do not discriminate between serum IgA (monomeric and dimeric) and higher molecular forms such as secretory IgA. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <u>Working dilutions:</u> For histochemical and cytochemical use are usually between 1/100 and 1/500. In ELISA and comparable non-precipitating antibody-binding assays between 1/1000 and 1/5000.
Reactivity:	Canine
Host:	Goat
Immunogen:	Purified normal IgA isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 No preservative added, as it may interfere with the antibody activity. No foreign proteins added. Label: Biotin State: Lyophilised hyperimmune Ig fraction Molar ratio: 6,9
Reconstitution Method:	Restore with 1 ml sterile distilled water



[View online »](#)

Concentration:	10 mg/ml
Purification:	DEAE-column Chromatography
Conjugation:	Biotin
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one week or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.