

Product datasheet for **AP31541AF-N**

Canine IgG (H+L chain) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Canine IgG (H+L chain) Goat Polyclonal Antibody
Applications:	ELISA, ID, IF, IP, WB
Recommended Dilution:	Can be used as unlabelled primary or secondary reagent for indirect detection techniques, to prepare conjugates with markers of the user's own choice, to prepare an insoluble immunoaffinity adsorbent or a solid phase antibody reagent by coupling to an artificial carrier and as catching or detection antibody in non-isotopic methodology and solid phase immunochemistry. When applied in any cytochemical or histochemical procedure or solids phase coupling technique, the optimum concentration of the IgG preparation should always be established by titration. <u>Recommended working dilutions:</u> In histochemistry are usually between 1/50 and 1/250. In ELISA and comparable non-precipitating antibody-binding assays between 1/500 and 1/5000.
Reactivity:	Canine
Host:	Goat
Immunogen:	Purified normal IgG isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 No preservative added, as it may interfere with the antibody activity. No foreign protein added. State: Azide Free State: Lyophilised hyperimmune Ig fraction
Reconstitution Method:	Restore with 1 ml sterile distilled water.
Concentration:	10 mg/ml
Purification:	DEAE-column Chromatography
Conjugation:	Unconjugated



[View online »](#)

Storage:

Prior to reconstitution store at 2-8°C.
Following reconstitution store undiluted at 2-8°C for one week
or (in aliquots) at -20°C for longer.
Avoid repeated freezing and thawing.