

Product datasheet for **AP31541HR-N**

Canine IgG (H+L chain) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Canine IgG (H+L chain) Goat Polyclonal Antibody
Applications:	ELISA, ID, IF, IHC, IP, WB
Recommended Dilution:	Can be used in enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG, antigen or antibody, of appropriately treated cell and tissue substrates at the cellular and subcellular level. In non-isotopic assay methodology (e.g. ELISA) to identify and measure a specific IgG in dog serum or other body fluid. In electron microscopy, since the complex between the conjugated antibody and the antigen also has electron-dense properties. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <u>Working dilutions:</u> For histochemical and cytochemical use are usually between 1/100 and 1/500. In ELISA and comparable non-precipitating antibody-binding assays between 1/1000 and 1/10000.
Reactivity:	Canine
Host:	Goat
Immunogen:	Purified normal IgG isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 No preservative added, as it may interfere with the antibody activity. No foreign proteins added. Label: HRP State: Lyophilised hyperimmune Ig fraction Label: Horseradish Peroxidase Molar ratio: 1,3
Reconstitution Method:	Restore with 1 ml sterile distilled water
Concentration:	10 mg/ml



[View online »](#)

Purification:	DEAE-column Chromatography
Conjugation:	HRP
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one week or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.