

Product datasheet for **AP31542BT-N**

Canine IgM (Fc specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Canine IgM (Fc specific) Goat Polyclonal Antibody
Applications:	ID, IP
Recommended Dilution:	Can be used in immunocytochemical and immunohistochemical use for the detection of IgM at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgM antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using an reference antibody of goat origin known to be of the IgM isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgM in dog serum or other body fluids. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <u>Working dilutions:</u> For histochemical and cytochemical use are usually between 1/100 and 1/250. In ELISA and comparable non-precipitating antibody-binding assays between 1/1000 and 1/4000.
Reactivity:	Canine
Host:	Goat
Immunogen:	Purified normal IgM isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG
Formulation:	PBS, pH 7.2 No preservative added, as it may interfere with the antibody activity. No foreign proteins added. Label: Biotin State: Lyophilised hyperimmune Ig fraction Molar ratio: 5,5
Reconstitution Method:	Restore with 1 ml sterile distilled water



[View online »](#)

Concentration: 10 mg/ml

Purification: DEAE-column Chromatography

Conjugation: Biotin

Storage: Prior to reconstitution store at 2-8°C.
Following reconstitution store undiluted at 2-8°C for one week
or (in aliquots) at -20°C for longer.
Avoid repeated freezing and thawing.

Note: **Adsorption:** Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.