

Product datasheet for AP31895PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Col3a1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IF, IHC

Recommended Dilution: Recommended for use in Immunohistochemistry on Frozen Sections and for

Immunostaining of cultured rat cells.

Suitable for **Dot-blotting** and **ELISA** on native Rat Collagen type I.

Recommended Dilutions: 1/20 for Immunohistochemical procedures if Peroxidase labeled secondary antibodies are applied or 1/10 if a FITC labeled secondary antibody is used.

Reactivity: Rat

Host: Rabbit
Clonality: Polyclonal

Immunogen: Native Rat Collagen type I.

Antibodies to Rat Collagen type I are raised in rabbits which are numerously immunized with extensively purified native collagen type I extracted from rat tail tendon into dilute acidic buffer after mild pepsin digestion. Pooled antisera are passed over DEAE-cellulose to produce IgG-enriched fraction. Next, the antisera fraction is absorbed with immunobilized rat collagen types III to remove cross-reactive antibodies to antigenic determinants common for this type of collagen. This affinity purified antibody is obtained by binding to immobilized native rat collagen type I (the antigen used for immunization), followed by elution with acidic buffer,

neutralisation, dialysis, dispensing and lyophilization.

Specificity: Specificity was ascertained by competition ELISA.

Complete inhibition was found if the antibody was preincubated with Rat Collagen type I. Inhibition by Rat Collagen type III is observed only at 10-20 times higher concentration.

No inhibition is found with Fibrinectin. Fibrogen and Laminin.

Characteristic immunostaining pictures of frozen sections of Rat kidney and skin are

produced to certify absence of crossreactivity with basement membranes.

Formulation: State: Aff - Purified

State: Lyophilized purified IgG fraction

Reconstitution Method: Restore with 1 ml distilled water and add preservative if preferred.

Concentration: lot specific





Col3a1 Rabbit Polyclonal Antibody - AP31895PU-N

Purification: Affinity Chromatography: This antibody is obtained by binding to immobilized native rat

Collagen type I (the antigen used for immunization), followed by elution with acidic buffer,

neutralisation, dialysis, dispensing and lyophilization.

Conjugation: Unconjugated

Gene Name: collagen, type III, alpha 1 Database Link:

Entrez Gene 84032 Rat

P13941

Background:

It is often extremely difficult to generate antibodies with specificities to collagens due to the uninterrupted "Glycine-X-Y" triplet repeat that is a necessary part of the triple helical structure. The development of type specific antibodies is dependent on NON-DENATURED three-dimensional epitopes - this may result in diminished reactivity of some antibodies with denatured collagen or formalin-fixed, paraffin embedded tissues. Anti-Collagen antibodies have been used for indirect trapping ELISA for quantitation of antigen in serum using a standard curve, for immunoprecipitation and for native (non-denaturing, non-dissociating)

PAGE and western blotting for highly sensitive qualitative analysis.

Synonyms: COL3A1