

Product datasheet for AP31985PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CSEN (KCNIP3) (N-term) Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: Peptide ELISA: 1/8000 (Detection limit).

Western blot: 1-3 μg/ml. In transfected HEK293 transiently expressing Human KCNIP3 a band

of approx. 30kDa was observed. No bands are observed in the non-transfected HEK293.

Reactivity: Bovine, Canine, Human

Host: Goat

Clonality: Polyclonal

Immunogen: Synthetic peptide with sequence from the N-terminus of Human KCNIP3 (NP_038462.1).

Specificity: This antibody is expected to recognize Isoform 1 of KCNIP3 (NP_038462.1) only.

Formulation: Tris saline, pH~7.3

State: Aff - Purified

State: Liquid purified Ig fraction

Stabilizer: 0.5% BSA

Preservative: 0.02% Sodium Azide

Concentration: lot specific

Purification: Purification: Affinity Chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 29.2 kDa according to NP 038462.1.

Database Link: Entrez Gene 30818 Human

Q9Y2W7






Background:

KChIP3 is a member of the family of voltage-gated potassium (Kv) channel-interacting proteins (KCNIPs), which belong to the recoverin branch of the EF-hand superfamily. Members of the KCNIP family are small calcium binding proteins. They all have EF-hand-like domains, and differ from each other in the N-terminus. They are integral subunit components of native Kv4 channel complexes. They may regulate A-type currents, and hence neuronal excitability, in response to changes in intracellular calcium. This protein is also shown to function as a calcium-regulated transcriptional repressor, and to interact with presenilins. Mutations in the presenilin genes have been implicated in Alzheimer's disease. Due to utilization of an alternate in-frame translation start codon, the gene for this protein encodes two isoforms with different sizes.

Synonyms:

KCNIP3, CSEN, DREAM, KCHIP3, DRE-antagonist modulator

Product images:



HEK293 lysate (10 ug protein in RIPA buffer) overexpressing Human KCNIP3 with C-terminal MYC tag probed with KCNIP3 antibody at 1 ug/ml in Lane A and probed with anti-MYC Tag (1/1000) in Lane C. Mock-transfected HEK293 probed with (1 mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.