

# Product datasheet for AP32808PU-N

## PRAK (MAPKAPK5) Goat Polyclonal Antibody

**Product data:** 

**Product Type: Primary Antibodies Applications:** ELISA, IHC, IP, WB

Recommended Dilution: Peptide ELISA: 1/32000 (Detection Limit).

> Western blot: 0.5-1 µg/ml. In transfected HEK293 transiently expressing either Mouse MK5 fused to GFP or the similar Mouse MK2 fused to GFP; only the expected band was observed

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

in the MK5 lane after AP32808PU-N labeling.

Data obtained from Dr. M. B. Menon, Inst. Biochemistry, Hannover Medical School, Germany. **Immunoprecipitation:** MK5/ERK3 double knockout Mouse Embryonic Fibroblasts (MEFs) retrovirally transduced wiith Mk5/Erk3 or empty vector (GFP) were lysed from confluent

plates and used for IP with 1.5 µg AP32808PU-N or AP32809PU-N.

Western blots of the Immunoprecipitation were labelled with Mouse anti-MK5 (SC) or with Rabbit anti-ERK3 serum (CST). An approx 55kDa of MK5 is only precipitated from lysates of those KO MEFs that have been rescued by the ERK3/MK5 expression constructas described in

PMID: 22508986.

Data obtained from Dr. M. B. Menon, Inst. Biochemistry, Hannover Medical School, Germany. Immunohistochemistry on Paraffin Sections: 4-8 µg/ml. In paraffin embedded Human

Kidney shows brushborder staining in the PCT.

Reactivity: Bovine, Canine, Human, Mouse, Porcine, Rat

Host: Goat

Clonality: Polyclonal

Immunogen: Peptide with sequence from the internal region of the protein sequence according to

NP 003659.2; NP 620777.1.

Specificity: Recognizes MK5 / MAPKAPK5.

Formulation: Tris saline, pH~7.3

State: Aff - Purified

State: Liquid purified Ig fraction

Stabilizer: 0.5% BSA

Preservative: 0.02% Sodium Azide

Concentration: lot specific





### PRAK (MAPKAPK5) Goat Polyclonal Antibody - AP32808PU-N

Purification: Ammonium Sulphate Precipitation followed by antigen Affinity Chromatography using the

immunizing peptide

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.


**Gene Name:** mitogen-activated protein kinase-activated protein kinase 5

Database Link: Entrez Gene 8550 Human


Q8IW41

**Synonyms:** MAPK-activated protein kinase 5, MAPKAP kinase 5, MAPKAPK5, PRAK

## **Product images:**




HEK293 lysate (10 ug protein in RIPA buffer) overexpressing Mouse MK5-GFP (first lane) or Mouse MK2-GFP (second lane) probed with (0.5 ug/ml) in right panel and with -N (0.5 ug/ml) on left panel, Primary incubations were for 2 hours. Detected by chemiluminescence.



(5 ug/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.





and (1.5 ug) immunoprecipitations from lysates of MK5/ERK3 double knockout MEFs, with (third and fifth lanes) and without (fourth and sixth lanes) rescued MK5/ERK3 expression through retroviral transduction. The corresponding lysates (first and second lane resp.) were analyzed in parallel in this Western blot labelled with mouse anti-MK5 / PRAK (and coprecipitation was measured using Rabbit anti-ERK3 in the lower panel).