

## Product datasheet for **AP32992AF-N**

### Mouse IgM (Fc specific) Rabbit Polyclonal Antibody

#### Product data:

**Product Type:** Secondary Antibodies

**Product Name:** Mouse IgM (Fc specific) Rabbit Polyclonal Antibody

**Applications:** ELISA, ID, IF, IP, WB

**Recommended Dilution:** Can be used:

As unlabelled primary or secondary antibody reagent for indirect detection of IgM in Mouse cells, tissues and body fluids in Immunofluorescence and Immunoenzyme assay methods.

For the production of immunoconjugates with a selected marker.

To prepare immunoaffinity adsorbents by coupling to an artificial carrier.

In non-isotopic methodology based on solid phase immunochemistry (e.g. ELISA), both as catching antibody and detection reagent.

In Western blotting.

When applied in any immunocytochemical or histochemical staining procedure or solid phase coupling techniques, the optimum concentration of this product should be established before.

*Recommended Working Dilutions:*

**Histochemistry:** 1/100-1/250.

**ELISA and comparable non-precipitating antibody-binding assays:** 1/1000-1/5000.

*Precipitin titre:* not less than 1/64 when tested against normal Mouse serum in agar-block immunodiffusion titration.

**Performance testing:** Titre, specificity and reactivity of the subclass specific cross-adsorbed IgG is further evaluated in a number of highly sensitive non-precipitating antibody-binding assay systems. These performance tests include direct single and double identification of cIg in mouse cell and tissue substrates and their evaluation as second antibody in the analysis of human cells and tissues after reacting with a primary monoclonal mouse antibody to a human antigen. The quantitative specific recognition ability is verified in double staining procedures together with reference reagents of known specificity and reactivity.

**Reactivity:** Mouse

**Host:** Rabbit

**Immunogen:** Highly purified homogenous IgM isolated from Mouse serum.  
Freund's complete adjuvant is used in the first step of the immunization procedure.



[View online »](#)

Isotype:	IgG
Formulation:	PBS, pH 7.2 State: Azide Free State: Lyophilized purified hyperimmune IgG fraction Stabilizer: None Preservative: None, as it may interfere with the antibody activity. No foreign protein added
Reconstitution Method:	Restore with 1 ml sterile distilled water
Concentration:	10 mg/ml
Purification:	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG fraction by DEAE Chromatography
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Note:	<b>Adsorption:</b> Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the elimination of antibodies to the common Ig/Fab portion. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.