

Product datasheet for **AP33082HR-N**

Sheep IgG (H+L chain) Rabbit Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Sheep IgG (H+L chain) Rabbit Polyclonal Antibody
Applications:	ELISA, IF, IHC, WB
Recommended Dilution:	ELISA.

Dot blot.

Immunoblotting.

Immunocytochemistry.

Immunohistochemistry Paraffin Sections.

Can be used in enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG, antigen or antibody, of appropriately treated cell and tissue substrates at the cellular and subcellular level. In non-isotopic assay methodology (e.g. ELISA) to identify and measure a specific IgG in sheep serum or other body fluid. In electron microscopy, since the complex between the conjugated antibody and the antigen also has electron-dense properties. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Recommended Working Dilutions:

Histochemistry and Cytochemical Use: 1/100-1/500.

ELISA and comparable non-precipitating antibody-binding assays: 1/1,000-1/10,000.

Reactivity:	Sheep
Host:	Rabbit
Immunogen:	Purified normal IgG isolated from pooled sheep serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Isotype:	IgG



[View online »](#)

Formulation:	PBS, pH 7.2 Label: HRP State: Lyophilized purified hyperimmune IgG fraction Stabilizer: None Preservative: None, as it may interfere with the antibody activity. Label: Horseradish peroxidase enriched for isoenzyme C (RZ = 3.2) Molar ratio: Enzyme/IgG ~1.7
Reconstitution Method:	Restore with 1 ml sterile distilled water.
Concentration:	10.0 mg/ml
Purification:	Hyperimmune antisera with strong activity are selected for fractionation by salt precipitation and purification of the IgG fraction by DEAE-chromatography.
Conjugation:	HRP
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Note:	Adsorption: Not required. Conjugation procedure: Conjugation is carried out using a proprietary modification of the periodate method, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. No foreign protein has been added. The conjugate is lyophilized to assure stability and long shelf life.