

Product datasheet for **AP33086HR-N**

Porcine IgG (Fc specific) Rabbit Polyclonal Antibody

Product data:

Product Type: Secondary Antibodies

Product Name: Porcine IgG (Fc specific) Rabbit Polyclonal Antibody

Applications: ELISA, IF, IHC, WB

Recommended Dilution: **ELISA.**

Dot blot.

Immunoblotting.

Immunocytochemistry.

Immunohistochemistry Paraffin Sections. Can be used in enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgG antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of swine origin known to be of the IgG isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgG in swine serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Recommended Working Dilutions:

Histochemistry and Cytochemical Use: 1/100-1/500.

ELISA and comparable non-precipitating antibody-binding assays: 1/1,000-1/10,000.

Reactivity: Porcine

Host: Rabbit

Immunogen: Purified normal IgG isolated from pooled Swine serum.
Freund's complete adjuvant is used in the first step of the immunization procedure.

Isotype: IgG



[View online »](#)

Formulation:	PBS, pH 7.2 Label: HRP State: Lyophilized purified hyperimmune IgG fraction Stabilizer: None Preservative: None, as it may interfere with the antibody activity. Label: Horseradish peroxidase enriched for isoenzyme C (RZ=3.2) Molar ratio: Peroxidase/IgG ~1.7
Reconstitution Method:	Restore with 1 ml sterile distilled water.
Concentration:	10.0 mg/ml
Purification:	Hyperimmune antisera with strong activity are selected for fractionation by salt precipitation and purification of the IgG fraction by DEAE-chromatography.
Conjugation:	HRP
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Note:	<p><u>Adsorption:</u> Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.</p> <p><u>Conjugation procedure:</u> Conjugation is carried out using a proprietary modification of the periodate technique for the binding to peroxidase, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilized to assure stability and long shelf life.</p>