

Product datasheet for **AP50751PU-N**

CCDC85C (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	FC, IHC, WB
Recommended Dilution:	ELISA: 1/1000. Western blot: 1/100 – 1/500. Immunohistochemistry on paraffin sections: 1/50 – 1/100. Flow Cytometry: 1/10 – 1/50.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	KLH conjugated synthetic peptide between 63–93 amino acids from the N-terminal region of human CC85C
Specificity:	This antibody reacts to CC85C.
Formulation:	PBS State: Aff – Purified State: Liquid purified Ig fraction Preservative: 0.09% (W/V) sodium azide
Concentration:	lot specific
Purification:	Affinity chromatography on Protein A
Conjugation:	Unconjugated
Storage:	Store undiluted at 2–8°C for one month or (in aliquots) at –20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	45210 Da




Gene Name: coiled-coil domain containing 85C


Database Link: [Entrez Gene 317762 Human A6NKD9](#)

Synonyms: Coiled-coil domain-containing protein 85C


Product images:



CC85C Antibody (N-term) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the CC85C antibody detected the CC85C protein (arrow).



CC85C Antibody (N-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CC85C Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



CC85C Antibody (N-term) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.