

Product datasheet for AP51152PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

CGBP (CXXC1) (Center) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: This antibody is suitable for **Western blottung**.

The suggested dilution is 1/100-1/500.

Reactivity: Human, Mouse

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide from the central region (between 325-354 aa) of Human

CpG-binding protein/CXXC1.

Specificity: This antibody detects CpG-binding protein/CXXC1 (Center).

Formulation: PBS with 0.09% (W/V) Sodium Azide.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Purified through a protein A column; followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: Homo sapiens CXXC finger protein 1 (CXXC1), transcript variant 1

Database Link: Entrez Gene 74322 MouseEntrez Gene 30827 Human

Q9P0U4



CGBP (CXXC1) (Center) Rabbit Polyclonal Antibody - AP51152PU-N

Background:


Human CpG binding protein (hCGBP) contains three cysteine-rich domains, two of which exhibit homology to the plant homeodomain finger domain and the third cysteine-rich domain conforms to the CXXC motif identified in DNA methyltransferase, human trithorax, and methyl-CpG binding domain protein 1. A fragment of hCGBP that contains the CXXC domain binds to an oligonucleotide probe containing a single CpG site, and this complex is disrupted by distinct oligonucleotide competitors that also contain a CpG motif(s). CBGP plays an important part in embryonic viability and peri-implantation development. Native hCGBP is detected as an 88-kDa protein by Western analysis and is ubiquitously expressed. CGBP trans-activates promoters that contain CpG motifs but not promoters in which the CpG is ablated suggesting that hCGBP is a transcriptional activator that recognizes unmethylated CpG dinucleotides.

Synonyms: CXXC1, CFP1, CGBP, PCCX1, PHF18

Note: Molecular Weight: 75712 Da

Protein Families: Druggable Genome, Transcription Factors

Product images:



Wwestern blot analysis in mouse liver tissue lysates (35ug/lane) using CpG-binding protein/CXXC1 antibody. This demonstrates the CXXC1 antibody detected the CXXC1 protein (arrow).