

### Product datasheet for AP51228PU-N

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

#### Deformed Epidermal Autoregulatory Factor 1 (DEAF1) (C-term) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100-1/500.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

**Immunogen:** KLH conjugated synthetic peptide between 473-502 amino acids from the C-terminal region

of human DEAF1

**Specificity:** This antibody recognizes Human DEAF1 (C-term).

Formulation: PBS

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.09% Sodium Azide

**Concentration:** lot specific

**Purification:** Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: DEAF1, transcription factor

Database Link: Entrez Gene 10522 Human

<u>075398</u>



# Deformed Epidermal Autoregulatory Factor 1 (DEAF1) (C-term) Rabbit Polyclonal Antibody – AP51228PU-N

Background:


Transcription factor that binds to sequence with multiple copies of 5'-TTC[CG]G-3' present in its own promoter and that of the HNRPA2B1 gene. Down-regulates transcription of these genes. Binds to the retinoic acid response element (RARE) 5'-AGGGTTCACCGAAAGTTCA-3'. Activates the proenkephalin gene independently of promoter binding, probably through protein-protein interaction. When secreted, behaves as an inhibitor of cell proliferation, by arresting cells in the G0 or G1 phase. Required for neural tube closure and skeletal patterning. Regulates epithelial cell proliferation and side-branching in the mammary gland. Controls the expression of peripheral tissue antigens in pancreatic lymph nodes. Isoform 1 displays greater transcriptional activity than isoform 4. Isoform 4 may inhibit transcriptional activity of isoform 1 by interacting with isoform 1 and retaining it in the cytoplasm.

Synonyms: SPN, ZMYND5, Suppressin, SPN, NUDR

Note: Molecular Weight: 59327 Da

**Protein Families:** Secreted Protein, Transcription Factors, Transmembrane

## **Product images:**



Western blot analysis of DEAF1 Antibody (C-term) in 293 cell line lysates (35ug/lane). This demonstrates the DEAF1 antibody detected the DEAF1 protein (arrow).