

Product datasheet for AP51506PU-N

Troduct dutasification / il 5 15001 0 1

FABP2 (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100-1/500.

Immunohistochemistry on Paraffin Sections: 1/10-1/50.

Reactivity: Human
Host: Rabbit
Isotype: Ig

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of

Human FABP2

Specificity: This antibody recognizes Human FABP2 (N-term).

Formulation: PBS containing 0.09% (W/V) Sodium Azide as preservative

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: fatty acid binding protein 2

Database Link: Entrez Gene 2169 Human

P12104



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


The intracellular fatty acid-binding proteins (FABPs) belong to a multigene family with nearly twenty identified members. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Intestinal fatty acid-binding protein 2 gene contains four exons and is an abundant cytosolic protein in small intestine epithelial cells. This gene has a polymorphism at codon 54 that identified an alanine-encoding allele and a threonine-encoding allele. Thr-54 protein is associated with increased fat oxidation and insulin resistance. [provided by RefSeq].

Synonyms: FABP-2, Fatty acid-binding protein, intestinal, I-FABP, FABPI


Note: Molecular Weight: 15207 Da

Protein Pathways: PPAR signaling pathway

Product images:



Western blot analysis of FABP2 (arrow) using FABP2 Antibody (N-term). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the FABP2 gene.



Formalin fixed and paraffin embedded human stomach tissue reacted with FABP2 Antibody (Nterm) followed by peroxidase conjugation of the secondary antibody and DAB staining.