OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for AP51694PU-N

FNIP2 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	FC, IHC, WB
Recommended Dilution:	**ELISA:** 1/1000. **Western Blot:** 1/100-1/500. **Flow Cytometry:** 1/10-1/50. **Immunohistochemistry on Paraffin Sections:** 1/50-1/100.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	KLH conjugated synthetic peptide between 783-812 amino acids from the C-terminal region of Human FNIP2
Specificity:	This antibody recognizes Human FNIP2 (C-term).
Formulation:	PBS containing 0.09% (W/V) Sodium Azide as preservative State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	lot specific
Purification:	Protein A column, followed by peptide affinity purification
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	folliculin interacting protein 2
Database Link:	Entrez Gene 57600 Human Q9P278
Synonyms:	FNIPL, KIAA1450, MAPO1, FNIP1-like protein, Folliculin-interacting protein 2
Note:	**Molecular Weight:** 122115 Da

View online »

Product images:



Western blot analysis of FNIP2 Antibody (C-term) in Hela cell line lysates (35ug/lane). This demonstrates the FNIP2 antibody detected the FNIP2 protein (arrow).



Immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle reacted with FNIP2 Antibody (C-term) followed by peroxidase conjugation of the secondary antibody and DAB staining.



Flow cytometric analysis of Hela cells using FNIP2 Antibody (C-term) (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.