

Product datasheet for AP52064PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Heme Oxygenase 1 (HMOX1) (Center) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, WB

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100-1/500. **Flow Cytometry:** 1/10-1/50.

Reactivity: Human
Host: Rabbit
Isotype: Ig

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 184-212 amino acids from the Central region of

Human Heme oxygenase 1 / HMOX1

Specificity: This antibody recognizes Human Heme oxygenase 1 / HMOX1 (Center).

Formulation: PBS containing 0.09% (W/V) Sodium Azide as preservative

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein A column, followed by peptide affinity purification

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: Homo sapiens heme oxygenase 1 (HMOX1)

Database Link: Entrez Gene 3162 Human

P09601





Background: Heme oxygenase, an essential enzyme in heme catabolism, cleaves heme to form biliverdin,

which is subsequently converted to bilirubin by biliverdin reductase, and carbon monoxide, a putative neurotransmitter. Heme oxygenase activity is induced by its substrate heme and by various nonheme substances. Heme oxygenase occurs as 2 isozymes, an inducible heme oxygenase-1 and a constitutive heme oxygenase-2. HMOX1 and HMOX2 belong to the heme


oxygenase family.

Synonyms: HO-1, HO, HO1


Note: Molecular Weight: 32819 Da

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Porphyrin and chlorophyll metabolism

Product images:



Western blot analysis of HMOX1 (arrow) using Heme oxygenase 1 / HMOX1 Antibody (Center). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the HMOX1 gene.



Flow cytometric analysis of A549 cells using Heme oxygenase 1 / HMOX1 Antibody (Center) (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.