

Product datasheet for AP52069PU-N

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

HNRNPCL1 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IHC, WB

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100-1/500. **Flow Cytometry:** 1/10-1/50

Immunohistochemistry on Paraffin Sections: 1/50-1/100.

Reactivity: Human
Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 259-289 amino acids from the C-terminal region

of human HNRNPCL1.

Specificity: This antibody recognizes Human HNRNPCL1 (C-term).

Formulation: PBS containing 0.09% (W/V) Sodium Azide as preservative

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein A column, followed by peptide affinity purification

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: heterogeneous nuclear ribonucleoprotein C-like 1

Database Link: Entrez Gene 343069 Human

O60812

Background: May play a role in nucleosome assembly by neutralizing basic proteins such as A and B core

hnRNPs (By similarity).


Synonyms: HNRPCL1, hnRNP C-like 1






Note: Molecular Weight: 32142 Da

Product images:




Western blot analysis of HNRNPCL1 Antibody (Cterm) in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the HNRNPCL1 antibody detected the HNRNPCL1 protein (arrow).



Immunohistochemistry analysis in formalin fixed and paraffin embedded human uterus tissue reacted with HNRNPCL1 Antibody (C-term) followed by peroxidase conjugation of the secondary antibody and DAB staining.

MDA-MB435



Flow cytometric analysis of MDA-MB435 cells using HNRNPCL1 Antibody (C-term) (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.