

Product datasheet for AP52276PU-N

KIAA0090 (EMC1) (C-term) Rabbit Polyclonal Antibody

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product data:

Product Type: Primary Antibodies

Applications: FC, IHC, WB

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100-1/500. Flow Cytometry: 1/10-1/50.

Immunohistochemistry on Paraffin Sections: 1/50-1/100.

Reactivity: Human, Mouse

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 873-902 amino acids from the C-terminal region

of human KIAA0090

Specificity: This antibody recognizes Human and Mouse KIAA0090 (C-term).

Formulation: PBS containing 0.09% (W/V) Sodium Azide as preservative

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein A column, followed by peptide affinity purification

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: ER membrane protein complex subunit 1

Database Link: Entrez Gene 23065 Human

Q8N766


Synonyms: 2700016F22Rik; C230096C10Rik; Kiaa0090; mKIAA0090; RGD1310427; RP23-371E13.1

Note: Molecular Weight: 111759 Da







Product images:



Western blot analysis of KIAA0090 Antibody (Cterm) in HepG2 cell line lysates (35ug/lane).




Western blot analysis of KIAA0090 Antibody (Cterm) in mouse spleen tissue lysates (35ug/lane).



Immunohistochemistry analysis in formalin fixed and paraffin embedded human testis carcinoma reacted with KIAA0090 Antibody (C-term), which was peroxidase conjugated to the secondary antibody and followed by DAB staining.





Flow cytometric analysis of HepG2 cells using KIAA0090 Antibody (C-term) (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.